

## Technical Brochure

- ✓ Foliar
- ✓ Furrow
- ✓ Fertigation



### Hi-2 Trace

A chelated blend of Zinc, Copper and Sulphur built for fast uptake in foliar sprays, fertigation and furrow injection

#### Benefits

- ✓ Essential elements for multiple metabolic improvements including protein synthesis, auxin production, chlorophyll formation and conversion of starches to sugars.
- ✓ Stimulate root activity and early plant vigour when applied in furrow
- ✓ Improves natural defence mechanisms to disease.
- ✓ Increased plant sugar production and transport

#### Typical Analysis

	G/L	%W/V	%W/W
Sulphur (S)	68	6.8	5.1
Zinc (Zn)	113	11.3	8.6
Copper (Cu)	23	2.3	1.7
pH	3.6 - 4.0		
SG	1.20 – 1.34		

#### Application Rates

Foliar Application	Rate/Ha	Water	Furrow Injection	Rate/Ha	Water
Broadacre	0.5 - 2L	50 -100L	Broadacre	2 - 5L	80L
Horticulture	0.5 - 4L	300-1500L	<b>Fertigation</b>	<b>Rate/Ha</b>	<b>Water</b>
Orchards (stone fruit excl.)	0.5 - 1L	1000L	Horticulture	5 - 10L	Variable
Viticulture	0.5 - 3L	300-1200L	Viticulture	5 - 10L	Variable
			Orchards	5 -10L	Variable

**Important:** (1) A jar test should be done, to assess compatibility with other liquids.  
(2) Do not foliar spray any crop during flowering.

**Packaging:** 20L, 200L and 1000L

Version Created: 26<sup>th</sup> September 2023